



2014 AP[®] CHEMISTRY FREE-RESPONSE QUESTIONS



3. A student is given a standard galvanic cell, represented above, that has a Cu electrode and a Sn electrode. As current flows through the cell, the student determines that the Cu electrode increases in mass and the Sn electrode decreases in mass.
- Identify the electrode at which oxidation is occurring. Explain your reasoning based on the student's observations.
 - As the mass of the Sn electrode decreases, where does the mass go?
 - In the expanded view of the center portion of the salt bridge shown in the diagram below, draw and label a particle view of what occurs in the salt bridge as the cell begins to operate. Omit solvent molecules and use arrows to show the movement of particles.



2014 AP[®] CHEMISTRY FREE-RESPONSE QUESTIONS

- (d) A nonstandard cell is made by replacing the 1.0 M solutions of $\text{Cu}(\text{NO}_3)_2$ and $\text{Sn}(\text{NO}_3)_2$ in the standard cell with 0.50 M solutions of $\text{Cu}(\text{NO}_3)_2$ and $\text{Sn}(\text{NO}_3)_2$. The volumes of solutions in the nonstandard cell are identical to those in the standard cell.
- Is the cell potential of the nonstandard cell greater than, less than, or equal to the cell potential of the standard cell? Justify your answer.
 - Both the standard and nonstandard cells can be used to power an electronic device. Would the nonstandard cell power the device for the same time, a longer time, or a shorter time as compared with the standard cell? Justify your answer.
- (e) In another experiment, the student places a new Sn electrode into a fresh solution of 1.0 M $\text{Cu}(\text{NO}_3)_2$.

Half-Reaction	E° (V)
$\text{Cu}^+ + e^- \rightarrow \text{Cu}(s)$	0.52
$\text{Cu}^{2+} + 2e^- \rightarrow \text{Cu}(s)$	0.34
$\text{Sn}^{4+} + 2e^- \rightarrow \text{Sn}^{2+}$	0.15
$\text{Sn}^{2+} + 2e^- \rightarrow \text{Sn}(s)$	-0.14

- Using information from the table above, write a net-ionic equation for the reaction between the Sn electrode and the $\text{Cu}(\text{NO}_3)_2$ solution that would be thermodynamically favorable. Justify that the reaction is thermodynamically favorable.
- Calculate the value of ΔG° for the reaction. Include units with your answer.

2015 AP[®] CHEMISTRY FREE-RESPONSE QUESTIONS

CHEMISTRY

Section II


7 Questions

Time—1 hour and 45 minutes

YOU MAY USE YOUR CALCULATOR FOR THIS SECTION.

Directions: Questions 1–3 are long free-response questions that require about 23 minutes each to answer and are worth 10 points each. Questions 4–7 are short free-response questions that require about 9 minutes each to answer and are worth 4 points each.

Write your response in the space provided following each question. Examples and equations may be included in your responses where appropriate. For calculations, clearly show the method used and the steps involved in arriving at your answers. You must show your work to receive credit for your answer. Pay attention to significant figures.




1. Metal-air cells are a relatively new type of portable energy source consisting of a metal anode, an alkaline electrolyte paste that contains water, and a porous cathode membrane that lets in oxygen from the air. A schematic of the cell is shown above. Reduction potentials for the cathode and three possible metal anodes are given in the table below.

Half Reaction	E at pH 11 and 298 K (V)
$\text{O}_2(g) + 2 \text{H}_2\text{O}(l) + 4 e^- \rightarrow 4 \text{OH}^-(aq)$	+0.34
$\text{ZnO}(s) + \text{H}_2\text{O}(l) + 2 e^- \rightarrow \text{Zn}(s) + 2 \text{OH}^-(aq)$	-1.31
$\text{Na}_2\text{O}(s) + \text{H}_2\text{O}(l) + 2 e^- \rightarrow 2 \text{Na}(s) + 2 \text{OH}^-(aq)$	-1.60
$\text{CaO}(s) + \text{H}_2\text{O}(l) + 2 e^- \rightarrow \text{Ca}(s) + 2 \text{OH}^-(aq)$	-2.78


2015 AP[®] CHEMISTRY FREE-RESPONSE QUESTIONS

- (a) Early forms of metal-air cells used zinc as the anode. Zinc oxide is produced as the cell operates according to the overall equation below.




- (i) Using the data in the table above, calculate the cell potential for the zinc-air cell.
- (ii) The electrolyte paste contains OH^- ions. On the diagram of the cell above, draw an arrow to indicate the direction of migration of OH^- ions through the electrolyte as the cell operates.
- (b) A fresh zinc-air cell is weighed on an analytical balance before being placed in a hearing aid for use.
- (i) As the cell operates, does the mass of the cell increase, decrease, or remain the same?
- (ii) Justify your answer to part (b)(i) in terms of the equation for the overall cell reaction.
- (c) The zinc-air cell is taken to the top of a mountain where the air pressure is lower.
- (i) Will the cell potential be higher, lower, or the same as the cell potential at the lower elevation?
- (ii) Justify your answer to part (c)(i) based on the equation for the overall cell reaction and the information above.
- (d) Metal-air cells need to be lightweight for many applications. In order to transfer more electrons with a smaller mass, Na and Ca are investigated as potential anodes. A 1.0 g anode of which of these metals would transfer more electrons, assuming that the anode is totally consumed during the lifetime of a cell? Justify your answer with calculations.
- (e) The only common oxide of zinc has the formula ZnO .
- (i) Write the electron configuration for a Zn atom in the ground state.
- (ii) From which sublevel are electrons removed when a Zn atom in the ground state is oxidized?

2016 AP[®] CHEMISTRY FREE-RESPONSE QUESTIONS



3. To determine the molar mass of an unknown metal, M, a student reacts iodine with an excess of the metal to form the water-soluble compound MI_2 , as represented by the equation above. The reaction proceeds until all of the I_2 is consumed. The $MI_2(aq)$ solution is quantitatively collected and heated to remove the water, and the product is dried and weighed to constant mass. The experimental steps are represented below, followed by a data table.



Data for Unknown Metal Lab	
Mass of beaker	125.457 g
Mass of beaker + metal M	126.549 g
Mass of beaker + metal M + I_2	127.570 g
Mass of MI_2 , first weighing	1.284 g
Mass of MI_2 , second weighing	1.284 g

- (a) Given that the metal M is in excess, calculate the number of moles of I_2 that reacted.
- (b) Calculate the molar mass of the unknown metal M.

The student hypothesizes that the compound formed in the synthesis reaction is ionic.

- (c) Propose an experimental test the student could perform that could be used to support the hypothesis. Explain how the results of the test would support the hypothesis if the substance was ionic.

2016 AP[®] CHEMISTRY FREE-RESPONSE QUESTIONS



The student hypothesizes that Br_2 will react with metal M more vigorously than I_2 did because Br_2 is a liquid at room temperature.

- (d) Explain why I_2 is a solid at room temperature whereas Br_2 is a liquid. Your explanation should clearly reference the types and relative strengths of the intermolecular forces present in each substance.

While cleaning up after the experiment, the student wishes to dispose of the unused solid I_2 in a responsible manner. The student decides to convert the solid I_2 to $\text{I}^-(aq)$ anion. The student has access to three solutions, $\text{H}_2\text{O}_2(aq)$, $\text{Na}_2\text{S}_2\text{O}_3(aq)$, and $\text{Na}_2\text{S}_4\text{O}_6(aq)$, and the standard reduction table shown below.

Half reaction	E° (V)
$\text{S}_4\text{O}_6^{2-}(aq) + 2 e^- \rightarrow 2 \text{S}_2\text{O}_3^{2-}(aq)$	0.08
$\text{I}_2(s) + 2 e^- \rightarrow 2 \text{I}^-(aq)$	0.54
$\text{O}_2(g) + 2 \text{H}^+(aq) + 2 e^- \rightarrow \text{H}_2\text{O}_2(aq)$	0.68



- (e) Which solution should the student add to $\text{I}_2(s)$ to reduce it to $\text{I}^-(aq)$? Circle your answer below. Justify your answer, including a calculation of E° for the overall reaction.



- (f) Write the balanced net-ionic equation for the reaction between I_2 and the solution you selected in part (e).

2017 AP® CHEMISTRY FREE-RESPONSE QUESTIONS

7. A student wants to determine the concentration of H_2O_2 in a solution of $\text{H}_2\text{O}_2(aq)$. The student can use one of two titrants, either dichromate ion, $\text{Cr}_2\text{O}_7^{2-}(aq)$, or cobalt(II) ion, $\text{Co}^{2+}(aq)$. The balanced chemical equations for the two titration reactions are shown below.



The half-reactions and the E° values for the systems related to the titrations above are given in the following table.


Half-Reaction	E° (V) at 298 K
$\text{Co}^{3+}(aq) + e^- \rightarrow \text{Co}^{2+}(aq)$	1.84
$\text{H}_2\text{O}_2(aq) + 2 \text{H}^+(aq) + 2 e^- \rightarrow 2 \text{H}_2\text{O}(l)$	1.77
$\text{Cr}_2\text{O}_7^{2-}(aq) + 14 \text{H}^+(aq) + 6 e^- \rightarrow 2 \text{Cr}^{3+}(aq) + 7 \text{H}_2\text{O}(l)$	1.33
$\text{O}_2(g) + 2 \text{H}^+(aq) + 2 e^- \rightarrow \text{H}_2\text{O}_2(aq)$	0.70

- (a) Use the information in the table to calculate the following.
- E° for the reaction between $\text{Cr}_2\text{O}_7^{2-}(aq)$ and $\text{H}_2\text{O}_2(aq)$ at 298 K
 - E° for the reaction between $\text{Co}^{2+}(aq)$ and $\text{H}_2\text{O}_2(aq)$ at 298 K
- (b) Based on the calculated values of E° , the student must choose the titrant for which the titration reaction is thermodynamically favorable at 298 K.
- Which titrant should the student choose? Explain your reasoning.
 - Calculate the value of ΔG° , in kJ/mol_{rxn} , for the reaction between the chosen titrant and $\text{H}_2\text{O}_2(aq)$.

STOP

END OF EXAM

2018 AP[®] CHEMISTRY FREE-RESPONSE QUESTIONS



6. A student sets up a galvanic cell at 298 K that has an electrode of $\text{Ag}(s)$ immersed in a 1.0 M solution of $\text{Ag}^+(aq)$ and an electrode of $\text{Cr}(s)$ immersed in a 1.0 M solution of $\text{Cr}^{3+}(aq)$, as shown in the diagram above.
- (a) The student measures the voltage of the cell shown above and discovers that it is zero. Identify the missing component of the cell, and explain its importance for obtaining a nonzero voltage.

Half-Reaction	E° (V)
$\text{Ag}^+(aq) + e^- \rightarrow \text{Ag}(s)$	+ 0.80
$\text{Cr}^{3+}(aq) + 3 e^- \rightarrow \text{Cr}(s)$?

- (b) The student adds the missing component to the cell and measures E°_{cell} to be +1.54 V. As the cell operates, Ag^+ ions are reduced. Use this information and the information in the table above to do the following.
- Calculate the value of E° for the half-reaction $\text{Cr}^{3+}(aq) + 3 e^- \rightarrow \text{Cr}(s)$.
 - Write the balanced net-ionic equation for the overall reaction that occurs as the cell operates.
 - Calculate the value of ΔG° for the overall cell reaction in $\text{J/mol}_{\text{rxn}}$.